

本製品は日本国法に基づき製造販売されたものです。また、この添付文書は日本国法に基づき作成した添付文書を翻訳したものであり、日本国外の法規に準じているわけではありません。

This product is manufactured and marketed for domestic sales in accordance with the Japanese laws.
This leaflet, translated from the original package insert created in accordance with Japanese laws, is not in compliance with laws outside Japan.

Designated 2nd Class OTC drug

Please read the package insert carefully before use and keep it with this medicine.

Therapeutic drug
for eczema/dermatitis

新リビメックスローション

NEW LIVIMEX KOWA LOTION

Precautions for use

To be avoided

(Failure to follow the directions below may worsen existing symptoms or increase the risk of adverse reactions)

1. Do not use:

Varicella (chickenpox), affected area of athletic foot/tinea, or of suppuration.

2. Do not apply to a large area on the face.

3. Do not use over a long period consecutively.

What to ask a doctor, pharmacist or registered salesperson

1. Ask a doctor, pharmacist or registered salesperson before use if you:

- (1) Are currently treated by a doctor.
- (2) Are pregnant or may possibly be pregnant.
- (3) Have ever had allergic reaction to any medicines, etc.
- (4) Have extensive lesions.
- (5) Have severe weeping or erosion.

2. The following symptoms have occurred due to this product, may be observed as adverse reactions. If any of the symptoms occur after use, stop use immediately and ask a doctor, pharmacist or registered salesperson with the package insert.

Related area	Symptom
Skin	Rash/redness, itching, feeling of tightness
Skin (affected area)	Dermatomycosis such as athletic foot/tinea, acne, purulent symptoms, persistent irritation

3. If symptoms do not improve after using for 5 to 6 days, stop use and ask a doctor, pharmacist or registered salesperson with the package insert.

Indications

Eczema, dermatitis, rash, itching, heat rash, insect bite, urticaria

Ingredients/contents (per gram)

● Prednisolone valerate acetate 3mg

[Inactive ingredients] Vaseline, liquid paraffin, cetanol, stearyl alcohol, carboxyvinyl polymer, sorbitan sesquioleate, polyoxyethylene hydrogenated castor oil, polysorbate 60, paraben, disodium edetate, citric acid, sodium hydroxide

Dosage/directions

Rub appropriate amount onto the affected areas several times a day.

<Precautions for dosage/directions>

- (1) Follow the dosage/directions.
- (2) This medicine should be applied to children only under supervision of their guardians.
- (3) Take care not to get it in your eyes. In case it gets into your eyes, immediately rinse them with water or lukewarm water.
In severe cases, immediately consult with an ophthalmologist.
- (4) For topical use only.
- (5) Shake thoroughly before use.
- (6) Do not wrap around the affected area with a poor air permeability material, such as a wrap film, where the medicine was just rubbed. Also, when applying to a part that comes in contact with diaper, do not use tight-fitting diapers or sealed pants made of vinyl.



(Continued overleaf)

Precautions for storage and handling

- (1) Store in a cool and dry place avoiding high temperatures, humidity, and direct sunlight, with cap tightly closed.
- (2) Keep out of reach of children.
- (3) Do not transfer the product to another container. (It may cause misuse and deterioration in the quality.)
- (4) Do not use this product after its validity date (printed on the outer box and on the label of container) has expired.

Marketing Authorization Holder Kowa Company, Ltd. 3-4-14, Nihonbashi-Honcho, Chuo-ku, Tokyo

Y1778 R1K A1