

本製品は日本国法に基づき製造販売されたものです。また、この添付文書は日本国法に基づき作成した添付文書を翻訳したものであり、日本国外の法規に準じているわけではありません。

This product is manufactured and marketed for domestic sales in accordance with the Japanese laws. This leaflet, translated from the original package insert created in accordance with Japanese laws, is not in compliance with laws outside Japan.

2nd Class OTC drug

Please read this outer box carefully before use and keep it with this medicine.

Gastrointestinal agent

キャベジンコーワαプラス

CABAGIN KOWA α PLUS

Precautions for use

To be avoided

(Failure to follow the directions below may worsen existing symptoms or increase the risk of adverse reactions)

1. Do not use the following medicines while taking this medicine.

Gastrointestinal analgesics and antispasmodics

2. Do not use during breastfeeding; avoid breastfeeding when using this medicine.

(An ingredient may pass into breast milk and cause rapid heartbeat in the infant.)

What to ask a doctor, pharmacist or registered salesperson

1. Ask a doctor, pharmacist or registered salesperson before use if you:

- (1) Are currently treated by a doctor.
- (2) Are pregnant or may possibly be pregnant.
- (3) Are elderly.
- (4) Have ever had allergic reaction to any medicines, etc.
- (5) Have the following symptoms:
Dysuria
- (6) Have been diagnosed with the following conditions:
Kidney disease, cardiac disease, glaucoma, thyroid dysfunction

2. The following symptoms, occurred due to this medicine, may be observed as adverse reactions. If any of the symptoms occur after use, stop use immediately and ask a doctor, pharmacist or registered salesperson with this outer box.

Related area	Symptom
Skin	Rash/redness, itching

3. The following symptom may occur after use; in case the symptom persists or worsens, stop use and ask a doctor, pharmacist or registered salesperson with this outer box.

Dry mouth

4. If symptoms do not improve after using for about two weeks, stop use and ask a doctor, pharmacist or registered salesperson with this outer box.

Other precautions

This product may reduce the flow of breast milk.

Indications

Stomach distress, gastrasthenia, heavy feeling in the stomach, stomachache, overeating, overdrinking, heartburn, nausea (retching, churning of stomach, nausea due to hangover/overdrinking, sickness, nausea), vomiting, loss of appetite, indigestion, gastric hyperacidity, belching, dysphagia, promoting digestion, feeling of fullness in the stomach/abdomen, and stomach heaviness.

Dosage/directions

Take the tablets with room temperature or warm water after meals.

Age	One dose	Dose frequency
Adults (15 years and older)	2 tablets ○○	3 times daily
8-14 years of age	1 tablet ○	
Children (Under 8 years of age)	✗ Do not use.	

<Precautions for dosage/directions>

- (1) Follow the dosage/directions.
- (2) This medicine should be given to children only under supervision of their guardians.

(Continued overleaf)

Ingredients/contents (in 6 tablets)

●Methylmethionine sulfonium chloride 150.0 mg ●Sodium bicarbonate 600.0 mg ●Magnesium hydroxide 210.0 mg
●Precipitated calcium carbonate 1340.0 mg ●Scopolia extract powder (33.3%) 90.0 mg (30.0 mg as scopolia extract)
●Perilla herb extract powder 20.0 mg (180.0 mg as perilla herb) ●Magnolia bark extract powder 15.0 mg (180.0 mg as magnolia bark) ●Powdered swertia herb 30.0 mg ●Biodiastase 2000 24.0 mg ●Lipase AP12 15.0 mg
[Inactive ingredients] Hydroxypropyl cellulose, hydrogenated oil, carmellose calcium, corn starch, magnesium stearate, glyceryl stearate, polyoxyl stearate, shellac, talc, cinnamon bark, (partially hydrolyzed) polyvinyl alcohol, cellulose, silicon dioxide, *l*-menthol, dextrin

<Precautions for ingredients/contents>

This product has a unique smell due to the ingredients such as methylmethionine sulfonium chloride.

Precautions for storage and handling

- (1) Store in a cool and dry place avoiding high temperatures, humidity, and direct sunlight, with cap tightly closed.
- (2) Keep out of reach of children.
- (3) Do not transfer the product to another container.(It may cause misuse and deterioration in the quality.)
- (4) When tablets get wet with water, the smell may become stronger, or the water may alter the ingredients. Therefore, avoid falling water drops on tablets or handling tablets with wet hands. If tablets accidentally become wet, discard the wet tablets.
- (5) The cushioning material in the bottle is for protection of tablets during transportation. Make sure to discard the material once the cap is opened.
- (6) If the cap of the bottle is not closed tightly, the quality may deteriorate due to moisture in the air. Therefore, close the cap tightly after each use.
- (7) Write down the date when the cap is opened in the column on the outer box and on the label.
- (8) Do not use this product after its validity date (printed on the outer box and on the label) has expired. For quality maintenance reasons, this product should be used within 6 months once the cap is opened.



Marketing Authorization Holder Kowa Company, Ltd. 3-4-14, Nihonbashi-Honcho, Chuo-ku, Tokyo

C3K A1