

本製品は日本国法に基づき製造販売されたものです。また、この添付文書は日本国法に基づき作成した添付文書を翻訳したものであり、日本国外の法規に準じているわけではありません。

This product is manufactured and marketed for domestic sales in accordance with the Japanese laws.
This leaflet, translated from the original package insert created in accordance with Japanese laws, is not in compliance with laws outside Japan.

Designated 2nd Class OTC drug

Please read the package insert carefully before use and keep it with this medicine.

Drug for
insect bite/itching

ウナコーエースプレミアムL
UNA KOWA ACE PREMIUM L

Long-term use of this drug may cause dilation of blood vessels in skin, leading to redness, or thinning of skin. As face is more likely to be affected by skin redness, be especially careful when applying the drug to face. Do not use this drug as a make-up base or after shaving. Avoid chronic use of this drug without careful consideration after the symptoms improve.

Precautions for use

To be avoided

(Failure to follow the directions below may worsen existing symptoms or increase the risk of adverse reactions)

1. Do not use:

- (1) Varicella (chickenpox), affected area of athletic foot/tinea, or of suppuration.
- (2) Around surface of a wound.
- (3) In eyes, around the eyes, mucosa, etc.

2. Do not apply to a large area on the face.

3. Do not use over a long period consecutively.

What to ask a doctor, pharmacist or registered salesperson

1. Ask a doctor, pharmacist or registered salesperson before use if you:

- (1) Are currently treated by a doctor.
- (2) Are pregnant or may possibly be pregnant.
- (3) Have ever had allergic reaction to any medicines, etc.
- (4) Have extensive lesions.
- (5) Have severe weeping or erosion.

2. The following symptoms have occurred due to this product, may be observed as adverse reactions. If any of the symptoms occur after use, stop use immediately and ask a doctor, pharmacist or registered salesperson with the package insert.

Related area	Symptom
Skin	Rash/redness, itching, swelling
Skin (affected area)	Dermatomycosis such as athletic foot/tinea, acne, purulent symptoms, persistent irritation

3. If symptoms do not improve after using for 5 to 6 days, stop use and ask a doctor, pharmacist or registered salesperson with the package insert.

Ingredients/contents (per gram)

- Prednisolone valerate acetate 3.0mg ● Diphenhydramine hydrochloride 20.0mg ● Glycol salicylate 10.0mg
● *l*-Menthol 35.0mg ● *dl*-Camphor 10.0mg

[Inactive ingredients] Disodium edetate, citric acid, benzyl alcohol, ethanol

Indications

Insect bite, itching, eczema, rash, dermatitis, heat rash, urticaria

Dosage/directions


Apply appropriate amount to the affected areas several times a day.

(Continued overleaf)

<Precautions for dosage/directions>

- (1) Follow the dosage/directions.
- (2) This medicine should be applied to children only under supervision of their guardians.
- (3) Take care not to get it in your eyes. In case it gets into your eyes, immediately rinse them with water or lukewarm water.
In severe cases, immediately consult with an ophthalmologist.
- (4) For topical use only.
- (5) Do not wrap around the affected area with a poor air permeability material, such as a wrap film, where the medicine was just applied. When applying to a part that comes in contact with diaper, do not use tight-fitting diapers or sealed pants made of vinyl. In addition, avoid direct close contact of the applied skins when the medicine is used for the area such as back side of the knees or inside the bend of the elbows (ex. sitting straight with the knee together and the legs folded under the body).

[How to use the container]



* Although a sponge material has excellent usability and free from easy breakage is used for the product, note that the sponge head may break if it is not saturated sufficiently with the liquid.

Precautions for storage and handling

- (1) Store in a cool place avoiding high temperatures and direct sunlight, with cap tightly closed.
- (2) Keep out of reach of children.
- (3) Do not transfer the product to another container. (It may cause misuse and deterioration in the quality.)
- (4) Do not touch the mucosa such as the eyes with hands exposed to this medicine.
- (5) Do not leave the product in any place where it is exposed to high temperatures, such as in a car, as it may cause deformation of a container. Note that deformation of the container may cause sponge head drop out or liquid leakage.
- (6) If the medicine adheres to clothes or bedclothes and they become soiled, wash them with water or a cleanser as quickly as possible.
- (7) Avoid contact of the medicine with metals such as glasses, watches, and accessories, clothes, plastics, the coated surface of the floor or furniture, because the adhesion may cause their deterioration.
- (8) Keep fire away from this product.
- (9) Do not use this product after its validity date (printed on the outer box and on the label of container) has expired.



- If you have symptoms such as extended edema or swelling, difficulty in breathing, nausea, diarrhea, and pyrexia after getting a bee sting, etc., seek medical advice immediately.
- Your condition may not subside with this drug when the symptoms are severe or the affected area has extended. In this case, seek medical advice.

Marketing Authorization Holder Kowa Company, Ltd. 3-4-14, Nihonbashi-Honcho, Chuo-ku, Tokyo

Y1159 M3K A1