

本製品は日本国法に基づき製造販売されたものです。また、この添付文書は日本国法に基づき作成した添付文書を翻訳したものであり、日本国外の法規に準じているわけではありません。

This product is manufactured and marketed for domestic sales in accordance with the Japanese laws.  
This leaflet, translated from the original package insert created in accordance with Japanese laws, is not in compliance with laws outside Japan.

Designated 2nd Class OTC drug

Please read the package insert carefully before use and keep it with this medicine.

Therapeutic drug  
for eczema/dermatitis

## 新リビメックス<sup>®</sup>Wクリーム NEW LIVIMEX KOWA W CREAM

Long-term use of this drug may cause dilation of blood vessels in skin, leading to redness, or thinning of skin. As face is more likely to be affected by skin redness, be especially careful when applying the drug to face. Do not use this drug as a make-up base or after shaving. Avoid chronic use of this drug without careful consideration after the symptoms improve.

### Precautions for use

#### To be avoided

(Failure to follow the directions below may worsen existing symptoms or increase the risk of adverse reactions)

**1. Do not use:**

Varicella (chickenpox), affected area of athletic foot/tinea, or of suppuration.

**2. Do not apply to a large area on the face.**

**3. Do not use over a long period consecutively.**

#### What to ask a doctor, pharmacist or registered salesperson

**1. Ask a doctor, pharmacist or registered salesperson before use if you:**

- (1) Are currently treated by a doctor.
- (2) Are pregnant or may possibly be pregnant.
- (3) Have ever had allergic reaction to any medicines, etc.
- (4) Have extensive lesions.
- (5) Have severe weeping or erosion.

**2. The following symptoms have occurred due to this product, may be observed as adverse reactions. If any of the symptoms occur after use, stop use immediately and ask a doctor, pharmacist or registered salesperson with the package insert.**

Related area	Symptom
Skin	Rash/redness, itching, swelling
Skin (affected area)	Dermatmycosis such as athletic foot/tinea, acne, purulent symptoms, persistent irritation

**3. If symptoms do not improve after using for 5 to 6 days, stop use and ask a doctor, pharmacist or registered salesperson with the package insert.**

### Indications

Eczema, dermatitis, rash, itching, heat rash, insect bite, urticaria

### Ingredients/contents (per gram)

● Prednisolone valerate acetate 3mg ● Diphenhydramine hydrochloride 20mg ● Allantoin 2mg  
● Tocopherol acetate 5mg

[Inactive ingredients] Liquid paraffin, macrogol, isostearyl glyceryl ether, polyoxyethylene behenyl ether, citric acid, sodium citrate, glycerin, magnesium sulfate, paraben, fragrance

### Dosage/directions

Rub appropriate amount onto the affected areas several times a day.

(Continued overleaf)

### <Precautions for dosage/directions>

- (1) Follow the dosage/directions.
- (2) This medicine should be applied to children only under supervision of their guardians.
- (3) Take care not to get it in your eyes. In case it gets into your eyes, immediately rinse them with water or lukewarm water.

In severe cases, immediately consult with an ophthalmologist.

- (4) For topical use only.
- (5) Do not wrap around the affected area with a poor air permeability material, such as a wrap film, where the medicine was just rubbed. Also, when applying to a part that comes in contact with diaper, do not use tight-fitting diapers or sealed pants made of vinyl.



#### [How to use the container]



Press the sharp point of the cap into the sealed end of the tube.  
Push down firmly until the seal is open.

- Be sure to screw the cap on tightly after use.
- Do not bend or fold the tube. (If the tube is bent or folded, this may cause breakage of the tube.)

### Precautions for storage and handling

- (1) Store in a cool and dry place avoiding high temperatures, humidity, and direct sunlight, with cap tightly closed.
- (2) Keep out of reach of children.
- (3) Do not transfer the product to another container. (It may cause misuse and deterioration in the quality.)
- (4) Do not use this product after its validity date (printed on the outer box and on the tube) has expired.

Marketing Authorization Holder Kowa Company, Ltd. 3-4-14, Nihonbashi-Honcho, Chuo-ku, Tokyo

Y2797 T4K A1